

Crossovers between Superconducting Symmetry Classes[†]

V. A. Koziy and M. A. Skvortsov

*Landau Institute for Theoretical Physics, Russian Academy of Sciences,
Chernogolovka, Moscow region, 142432 Russia*

*Moscow Institute of Physics and Technology,
Dolgoprudnyi, Moscow region, 141700 Russia*

Received June 20, 2011

We study the average density of states in a small metallic grain coupled to two superconductors with the phase difference π , in a magnetic field. The spectrum of the low-energy excitations in the grain is described by the random matrix theory whose symmetry depends on the magnetic field strength and coupling to the superconductors. In the limiting cases, a pure superconducting symmetry class is realized. For intermediate magnetic fields or couplings to the superconductors, the system experiences a crossover between different symmetry classes. With the help of the supersymmetric σ -model we derive the exact expressions for the average density of states in the crossovers between the symmetry classes A–C and CI–C.

DOI: 10.1134/S0021364011150057

INTRODUCTION

Energy levels in small metallic particles with chaotic electron dynamics are random numbers. It is generally accepted that their spectral statistics in the ergodic regime is described by the random matrix theory (RMT) [1]. For disordered grains, this had been proved by Efetov [2] with the help of the supersymmetry technique [3], while for quantum billiards in the absence of disorder this statement is usually referred to as the Bohigas conjecture [4].

In the RMT, a system is characterized solely by its symmetry. In the application to condensed matter, the standard three Wigner–Dyson ensembles (orthogonal, unitary and symplectic) [5] describe level statistics in small metallic grains in the presence or absence of the time-reversal and spin-rotation symmetries [3].

Recently, the Wigner–Dyson classification had been extended to superconducting [6] and chiral [7] symmetry classes, which arise when the Hamiltonian possesses an additional symmetry with respect to changing the sign of the energy (counted from the Fermi energy). With the appearance of a selected energy point, in the superconducting/chiral classes even the average density of states (DOS), $\langle \rho(E) \rangle$, becomes a nontrivial function of the energy. This should be contrasted to the standard Wigner–Dyson ensembles where $\langle \rho(E) \rangle = \delta^{-1} = \text{const}$ and the first nontrivial quantity is the pair correlation function $R_2(\omega) = \delta^2 \langle \rho(E + \omega)\rho(E) \rangle - 1$.

The symmetry classes (three Wigner–Dyson, four superconducting and three chiral) correspond to the limits when various symmetries are either present or

completely broken. In the intermediate cases, the system experiences a crossover between different symmetry classes. The pair correlation function in the crossover between the orthogonal and unitary classes was obtained in [8, 9]:

$$R_2^{O-U}(\omega) = 1 - \frac{\sin^2 x}{x^2} + \int_1^\infty d\lambda \frac{\sin \lambda x}{\lambda} e^{-\alpha \lambda^2} \int_0^1 d\mu \mu \sin \mu x e^{\alpha \mu^2}, \quad (1)$$

where $x = \pi\omega/\delta$ and α is the symmetry-breaking parameter. Equation (1) interpolates between the orthogonal ($\alpha = 0$) and unitary ($\alpha = \infty$) results. The pair correlator is known also in the symplectic–unitary crossover [10, 11], its form being similar to Eq. (1).

The purpose of this Letter is to theoretically study crossovers between superconducting classes.

We will calculate the average density of states in a small diffusive metallic grain coupled to two superconducting terminals through tunnel barriers, see Fig. 1. The terminals have the phase difference π ensuring the absence of the minigap in the excitation spectrum [12]. A magnetic field H is applied to the system. The spin-rotation symmetry is assumed to be intact. We will be interested in the ergodic regime, $E \ll E_{\text{Th}}$, where $E_{\text{Th}} = D/L^2$ is the Thouless energy, D is the diffusion constant, and L is the grain size.

Under these conditions, the excitation spectrum in the grain can be described in terms of the RMT in the crossover region between the four symmetry classes shown in Fig. 2.

[†]The article is published in the original.




Fig. 1. Normal-metal dot coupled to two superconducting terminals with the phase difference π , in a magnetic field. NS interfaces are characterized by the set of transparencies $\{T_i\}$.

MAPPING TO EFETOV'S σ -MODEL

First attempts of field-theoretical description of hybrid NS systems [13–17] inspired by the identification of superconducting symmetry classes [6] have used the Bogoliubov–de Gennes (BdG) Hamiltonian as the starting point,

$$\hat{\mathcal{H}}_{\text{BdG}} = \begin{pmatrix} H & \Delta \\ \Delta^* & -H^T \end{pmatrix}, \quad (2)$$

where H is the single-particle Hamiltonian, and $\Delta(\mathbf{r})$ is the pairing field. The average quasiparticle DOS,

$$\langle \rho(E, \mathbf{r}) \rangle = -\text{Im} \langle \text{tr} \mathcal{G}_E^{\text{R}}(\mathbf{r}) \rangle / \pi, \quad (3)$$

is expressed in terms of the retarded Green's function of the BdG Hamiltonian,

$$\mathcal{G}_E^{\text{R}} = (E - \hat{\mathcal{H}}_{\text{BdG}} + i0)^{-1}, \quad (4)$$

which is then represented as a functional integral over an 8×8 supermatrix field Q acting in the direct product $\text{FB} \otimes \text{N} \otimes \text{PH}$ of the Fermi–Bose (FB), Nambu (N), and Particle–Hole (PH) spaces (spin-symmetric case is considered).

In hybrid NS systems, Andreev reflection off the order parameter field Δ couples the states with opposite energies, E and $-E$. So, the Nambu–Gor'kov Green's function \mathcal{G}_E^{R} essentially involves a pair of the retarded and advanced normal-metal Green's functions, G_E^{R} and G_{-E}^{A} . In the absence of the superconducting pairing field, $\Delta(\mathbf{r})$, correlations between the latter are conveniently described by Efetov's supersymmetric σ -model [2, 3] of the orthogonal symmetry class, with an 8×8 superfield Q acting in the direct product $\text{FB} \otimes \text{RA} \otimes \text{TR}$ of the Fermi–Bose (FB), Retarded–Advanced (RA) and Time–Reversal (TR) spaces (again we assume no spin interactions).

Thus, in studying the proximity effect in the normal part of a hybrid system, it is tempting to reformulate the field theory of [13–17] in the language of Efetov's supersymmetric σ -model. Provided that the inverse proximity effect in the superconducting




Fig. 2. Crossovers between spin-symmetric symmetry classes driven by the magnetic field (H) and coupling to superconductors (Δ). The dimensions of the FF- and BB-sectors of the supersymmetric σ -model for the average density of states are shown by $(n_{\text{F}}, n_{\text{B}})$.

regions can be neglected (rigid boundary conditions), Andreev scattering of normal electrons off the superconducting terminal will be viewed as an effective boundary condition at the NS interface mixing the R and A components of the field Q . Such a description is close in spirit to the scattering approach [18].

The average local DOS is given by the functional integral over the normal-metal region [17]:

$$\langle \rho(E, \mathbf{r}) \rangle = \frac{\nu}{4} \text{Re} \int \text{str}(k\Lambda Q) e^{-S_{\text{D}}[Q] - S_{\text{T}}[Q]} DQ(\mathbf{r}), \quad (5)$$

where $S_{\text{D}}[Q]$ is the bulk action:

$$S_{\text{D}} = \frac{\pi\nu}{8} \int d\mathbf{r} \text{str} \{ D(\nabla Q + ie\mathbf{A}[\tau_3, Q])^2 + 4iE\Lambda Q \}, \quad (6)$$

and the action $S_{\text{T}}[Q] = S_{\Gamma_1}[Q] + S_{\Gamma_2}[Q]$ describes NS interfaces [3, 19]:

$$S_{\Gamma_a} = -\frac{1}{2} \sum_i \text{str} \ln [1 + e^{-2\beta_i} Q_S^{(a)} Q^{(a)}]. \quad (7)$$

Here, ν is the DOS per one spin projection at the Fermi level, \mathbf{A} is the vector potential, $Q^{(a)}$ labels the Q field at the boundary with the a th superconductor, and the NS interface is specified by transmission coefficients $T_i = 1/\cosh^2 \beta_i$, with i labeling open channels. The field Q satisfies $Q^2 = 1$ and is subject to an additional symmetry constraint

$$Q = C Q^{\text{T}} C^{\text{T}}. \quad (8)$$

In the NS σ -model for $\langle \mathcal{G}_E^{\text{R}} \rangle$ [17], the matrices Λ , τ_3 , and C are given by the first column of table. An

Basic matrices in the two versions of the σ -model

	NS σ -model [17]	Efetov's σ -model [3]
Space	FB \otimes N \otimes PH	FB \otimes RA \otimes TR
Λ	$\sigma_z^N \sigma_z^{\text{PH}}$	σ_z^{RA}
τ_3	σ_z^N	σ_z^{TR}
C	$-\sigma_x^N \begin{pmatrix} i\sigma_y^{\text{PH}} & 0 \\ 0 & \sigma_x^{\text{PH}} \end{pmatrix}_{\text{FB}}$	$\sigma_z^{\text{RA}} \begin{pmatrix} i\sigma_y^{\text{TR}} & 0 \\ 0 & \sigma_x^{\text{TR}} \end{pmatrix}_{\text{FB}}$
$\hat{\Sigma}$	σ_x^N	$\begin{pmatrix} \sigma_1^{\text{RA}} \sigma_1^{\text{TR}} & 0 \\ 0 & \sigma_2^{\text{RA}} \sigma_2^{\text{TR}} \end{pmatrix}_{\text{FB}}$

exact mapping to Efetov's σ -model is realized by the similarity transformation $Q \mapsto VQV^{-1}$ with the matrix

$$V = \begin{pmatrix} -1_{\text{FB}} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1_{\text{FB}} \\ 0 & 1_{\text{FB}} & 0 & 0 \\ 0 & 0 & -k_{\text{FB}} & 0 \end{pmatrix}, \quad (9)$$

where the inner (outer) grading corresponds to the PH (N) space, and $k = \text{diag}(1, -1)_{\text{FB}}$ (we follow notations of [3]). Conjugation by V simultaneously transforms the matrices Λ , τ_3 , and C from NS representation to Efetov's representation given by the last column of table. This provides an exact mapping between the NS σ -model for Green's function of the BdG Hamiltonian, $\langle \mathcal{G}_E^{\text{R}} \rangle$, the standard Efetov's orthogonal σ -model for the product $\langle G_E^{\text{R}} G_{-E}^{\text{A}} \rangle$. On such a mapping both the structure of the manifold and the σ -model action get reproduced. We emphasize that this mapping takes place only in the normal part of a hybrid NS system, where the pairing amplitude $\Delta = 0$.

To complete the formulation of the model we have to specify the Q matrix in the bulk of a superconductor, Q_S . It has a familiar form parameterized with the help of the spectral angle $\theta_S = \arctan(i\Delta/E)$ as

$$Q_S = \Lambda \cos \theta_S + \hat{\Sigma} \sin \theta_S. \quad (10)$$

The most nontrivial ingredient of the mapping from the SN σ -model to Efetov's σ -model is the form of the matrix $\hat{\Sigma}$. In the initial NS representation [17] it is just

the Pauli matrix in the Nambu space: σ_x^{N} . Conjugating by V we get it in Efetov's representation:

$$\hat{\Sigma} = \begin{pmatrix} 0 & \Sigma \\ \Sigma^{-1} & 0 \end{pmatrix}_{\text{TR}}, \quad \Sigma = \begin{pmatrix} 0 & k_{\text{FB}} \\ 1_{\text{FB}} & 0 \end{pmatrix}_{\text{RA}}. \quad (11)$$

We see that superconducting boundary conditions "violate supersymmetry": the matrix $\hat{\Sigma}$ acts as $\sigma_1^{\text{RA}} \sigma_1^{\text{TR}}$ in the FF-block and as $\sigma_2^{\text{RA}} \sigma_2^{\text{TR}}$ in the BB-block. This is the reason why a nontrivial DOS can be obtained by integration (5) over the standard orthogonal σ -model manifold.

ZERO-DIMENSIONAL LIMIT

In the ergodic regime, $E \ll E_{\text{Th}}$, the functional integral (5) is dominated by the zero mode, $Q(\mathbf{r}) = \text{const}$. We will be interested in the average global DOS normalized by the inverse mean quasiparticle level spacing $\delta = (2\nu V)^{-1}$:

$$\langle \rho(E) \rangle = \delta \int \langle \rho(E, \mathbf{r}) \rangle d\mathbf{r}. \quad (12)$$

This quantity can be written as an integral over a single 8×8 supermatrix Q :

$$\langle \rho(E) \rangle = \frac{1}{8} \text{Re} \int \text{str}(k\Lambda Q) e^{-S[Q]} DQ, \quad (13)$$

with the action consisting of three terms:

$$S[Q] = \frac{ix}{4} \text{str} \Lambda Q - \frac{\alpha}{4} \text{str}(\tau_3 Q)^2 + \frac{\gamma}{8} \text{str}(\hat{\Sigma} Q)^2. \quad (14)$$

Here, $x = \pi E/\delta$, and the symmetry breaking parameters α and γ are given by

$$\alpha = \pi \nu D e^2 \int \mathbf{A}^2 d\mathbf{r}, \quad \gamma = \frac{G_A}{8}. \quad (15)$$

One can estimate $\alpha \sim (\phi/\phi_0)^2/g$, where ϕ is the flux through the grain, ϕ_0 is the flux quantum, and $g \sim E_{\text{Th}}/\delta \gg 1$ is the dimensionless grain conductance. The last term in the action (14) is written in the tunneling limit, $T_i \ll 1$, and the parameter γ is expressed through the dimensionless (in units of e^2/h) Andreev conductance of the grain [20], $G_A = 2 \sum_i T_i^2$ (the factor 2 accounts for two NS interfaces assumed to be identical, see Fig. 1). A strong magnetic field randomizes electron phase and the crossover from the unitary class to class C can be obtained with just one superconducting terminal attached to the grain (in that case $G_A = \sum_i T_i^2$). The last term in Eq. (14) is written in the subgap limit, $E \ll \Delta$, when $Q_S = \hat{\Sigma}$.

Equations (13) and (14) describe the average DOS in the two-parametric crossover between the four sym-

metry classes shown in Fig. 2. Instead of studying the complicated general behavior, we will restrict ourselves to two one-parametric crossovers: class A–class C ($\alpha = \infty$, γ arbitrary) and class CI–class C ($\gamma = \infty$, α arbitrary).

CLASS A–CLASS C CROSSOVER

In sufficiently strong magnetic fields ($\alpha \gg 1$), cooperon degrees of freedom get frozen and the Q matrix becomes diagonal in the TR-space:

$$Q = \begin{pmatrix} Q_u & 0 \\ 0 & k\Lambda Q_u^T \Lambda k \end{pmatrix}_{\text{TR}}, \quad (16)$$

where the 4×4 matrix $Q_u \in \text{FB} \otimes \text{RA}$ spans the manifold of the unitary Efetov’s σ -model. Therefore one can simply take the well-known Efetov’s parameterization of this manifold [3], in which the FF- and BB-sectors are parameterized by a radial variable ($\lambda_{\text{F,B}}$) and an angular variable ($\varphi_{\text{F,B}}$), with $-1 \leq \lambda_{\text{F}} \leq 1$ and $\lambda_{\text{B}} \geq 1$. It can be easily seen that the term $(\gamma/8)\text{str}(\hat{\Sigma} Q)^2$ in the action (14) does not depend on λ_{F} . Thus coupling to a superconductor suppresses only the BB degrees of freedom, shrinking it to a point at $\gamma \rightarrow \infty$ (see Fig. 2). Calculation of the integral (13) is straightforward leading to the exact expression for the class A–class C crossover:

$$\langle \rho(E) \rangle = 1 - 2\gamma \frac{\sin x}{x} \int_1^\infty d\lambda \lambda \cos \lambda x e^{-\gamma(\lambda^2 - 1)}. \quad (17)$$

The function $\langle \rho(E) \rangle$ is plotted in Fig. 3 for several values of the symmetry breaking parameter γ .

In the limit $\gamma \rightarrow \infty$, one recovers the C-class result:

$$\langle \rho_{\text{C}}(E) \rangle = 1 - \frac{\sin 2x}{2x}. \quad (18)$$

In the limit of weak coupling to a superconductor, $\gamma \ll 1$, the uniform metallic average DOS is perturbed in a small vicinity of the Fermi energy, at $E \lesssim \delta \sqrt{\gamma}$, but this perturbation is strong, completely suppressing the DOS at $E = 0$:

$$\langle \rho(E) \rangle \approx f\left(\frac{x}{2\sqrt{\gamma}}\right), \quad f(z) = 2ze^{-z^2} \int_0^z e^{t^2} dt. \quad (19)$$

Vanishing of $\langle \rho(0) \rangle$ can be explained by repulsion of energy levels E and $-E$ which becomes effective at very small energies $E \lesssim \delta \sqrt{\gamma}$. (Formally, noncommutativity of the limits $E \rightarrow 0$ and $\gamma \rightarrow 0$ is a consequence of the noncompactness of the BB-sector of the theory.)




Fig. 3. Average density of states, $\langle \rho(E) \rangle$, in the class A (unitary)–class C crossover. The curves correspond to different values of the symmetry-breaking parameter: $\gamma = 0$ (unperturbed DOS, $\langle \rho(E) \rangle = 1$), 0.01, 0.1, 1, and ∞ (class C).

Level statistics in classes A and C is known to be described by free fermions: The joint probability density of energy levels can be interpreted as the square of the ground-state wave function for a system of noninteracting one-dimensional (1D) fermions. Class A (unitary) corresponds to free fermions on a line [1], while class C corresponds to free fermions with the Dirichlet boundary condition at the origin [6]. Our result (17) indicates that in the crossover A–C energy levels cannot be considered as noninteracting fermions in an appropriate single-particle potential.

CLASS CI–CLASS C CROSSOVER

Now we turn to the case of strong coupling to a superconductor, $\gamma \gg 1$. The corresponding term in the action (14) enforces $\text{str}(\hat{\Sigma} Q)^2 = 0$. The latter condition together with the relation $\{\Lambda, \hat{\Sigma}\} = 0$ leads to the linear constraint

$$\{Q, \hat{\Sigma}\} = 0. \quad (20)$$

The next step in constructing the parameterization suitable for calculation in the crossover region is to study the commuting generators W of the CI-class manifold. Writing $Q = \Lambda(1 + W + \dots)$ with $\{W, \Lambda\} = 0$, and solving the linearized constraints (8) and (20), we find four commuting generators, three residing in the FF-sector, and one in the BB-sector (see Fig. 2):

$$W_{\text{com}} = \begin{pmatrix} W_{\text{FF}} & 0 \\ 0 & W_{\text{BB}} \end{pmatrix} = W_{\text{com}}^{(+)} + W_{\text{com}}^{(-)}, \quad (21)$$

where

$$W_{\text{FF}} = \frac{i}{2} \begin{pmatrix} 0 & 0 & z & -c \\ 0 & 0 & c & z^* \\ z^* & c & 0 & 0 \\ -c & z & 0 & 0 \end{pmatrix}_{\text{RA}}, \quad (22)$$

$$W_{\text{BB}} = \frac{q}{2} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}_{\text{RA}}, \quad (23)$$

with complex z and real c and q .

As the α -dependent term in the action (14) contains the symmetry-breaking matrix τ_3 , it is convenient to split W_{com} into a part $W_{\text{com}}^{(+)}$ commuting with τ_3 (z - and z^* -modes), and a part $W_{\text{com}}^{(-)}$ anticommuting with τ_3 (c - and q -modes). At finite α , the latter modes acquire a mass proportional to α . They will completely freeze out in the C-class limit ($\alpha \rightarrow \infty$), where only the modes $z = \theta e^{i\varphi}$ and $z^* = \theta e^{-i\varphi}$ will be unaffected, generating the sphere S^2 in the FF-sector.

In constructing the global parameterization of the CI-class manifold we will follow an approach of [9] in order to maximally simplify the symmetry breaking term $S_\alpha[Q] = -(\alpha/4)\text{str}(\tau_3 Q)^2$. We start with parameterizing the commuting content of Q as $Q_{\text{com}} = U_C^{-1} U_m^{-1} \Lambda U_m U_C$, where

$$U_C = \exp W_{\text{com}}^{(+)}, \quad U_m = \exp W_{\text{com}}^{(-)}. \quad (24)$$

With such a choice, $S_\alpha[Q_{\text{com}}]$ will explicitly depend only on the massive modes c and q .

Now we turn to the Grassmann content of the parameterization. We search for Grassmann generators \tilde{W} which obey $[\tau_3, \tilde{W}] = 0$ (to simplify the term S_α) and $[\Lambda, \tilde{W}] = 0$ (to simplify the term $\text{str} \Lambda Q$). Employing also linearized Eqs. (8) and (20), we find that these anticommuting generator are parameterized by two Grassmann numbers:

$$W_{\text{Gr}}^{(+)}[\xi, \rho] = \begin{pmatrix} u & 0 & 0 & 0 \\ 0 & v & 0 & 0 \\ 0 & 0 & v & 0 \\ 0 & 0 & 0 & -u \end{pmatrix}, \quad (25)$$

$$u = \begin{pmatrix} 0 & \xi \\ -\rho & 0 \end{pmatrix}_{\text{FB}}, \quad v = \begin{pmatrix} 0 & \rho \\ \xi & 0 \end{pmatrix}_{\text{FB}}. \quad (26)$$

The desired parameterization of the Q manifold in the spirit of [9] has the form

$$Q = U_\xi^{-1} U_C^{-1} U_\mu^{-1} U_m^{-1} \Lambda U_m U_\mu U_C U_\xi, \quad (27)$$

where the matrices U_m and U_C are defined in Eq. (24), and

$$U_\xi = \exp W_{\text{Gr}}^{(+)}[\xi, \rho], \quad (28)$$

$$U_\mu = \exp W_{\text{Gr}}^{(+)}[\mu, \eta].$$

After some algebra we obtain the Berezinian of the parameterization (27):

$$J = \frac{\sin \theta}{2(1 - \cos \theta)} \frac{\cos^2 c}{(\sin c + i \sinh q)^2} \equiv J_C J_m. \quad (29)$$

Similar to [9], it splits into factors depending either on massless or on massive coordinates.

The ingredients of the action (14) take a simple form:

$$\begin{aligned} \text{str}(\tau_3 Q)^2 &= 4(\cos 2c - \cosh 2q), \\ \text{str} \Lambda Q &= -4[P + (1 - \lambda)R\xi\rho], \end{aligned}$$

where $\lambda = \cos \theta$, $P = \cosh q - \lambda \cos c$, $R = \cosh q - \cos c$, while the pre-exponential factor in (13) is the most involved:

$$\begin{aligned} \text{str}(k \Lambda Q) &= 4[(\cosh q + \lambda \cos c) + 2P\eta\mu \\ &+ (1 + \lambda)R\xi\rho + 2(1 - \lambda)R\eta\mu\xi\rho + \dots], \end{aligned}$$

where the omitted terms do not contribute to the average DOS.

Let us expand the integrand in Eq. (13) in Grassmann variables:

$$F_{00} + F_{20}\xi\rho + F_{02}\mu\eta + F_{22}\xi\rho\mu\eta + \dots$$

The parameterization (27) is singular at $\theta = 0$ ($U_C = 1$) and at $c = q = 0$ ($U_m = 1$). Therefore the integral (13) will contain not only the regular contribution from the term F_{22} , but also the contributions from the terms F_{00} , F_{20} , and F_{02} , which are finite due to the Parisi–Sourlas–Efetov–Wegner theorem [21], in complete analogy with the calculation of [9]. In particular, the terms with F_{00} and F_{20} reproduce the C-class result (18), while the other two terms are responsible for the crossover. After some algebra, the general expression for the average DOS in the class CI–class C crossover takes the form

$$\begin{aligned} \langle \varrho(E) \rangle &= 1 - \frac{\sin 2x}{2x} \\ &+ \frac{1}{2\pi} \int_{-\infty}^{\infty} dq \int_{-\pi/2}^{\pi/2} dc \cos c \frac{\sinh^2 q - \sin^2 c}{\sinh^2 q + \sin^2 c} \quad (30) \end{aligned}$$

$$\times \sin(x \cos c) \sin(x \cosh q) \exp[\alpha(\cos 2c - \cosh 2q)].$$




Fig. 4. Average density of states, $\langle \rho(E) \rangle$, in the class CI–class C crossover. The curves correspond to different values of the symmetry-breaking parameter: $\alpha = 0$ (class CI, tiny oscillations), 0.2, 1, and ∞ (class C, large oscillations). Inset: $\langle \rho(E) \rangle$ for small deviations from class CI: $\alpha = 0, 0.01, 0.025, \text{ and } 0.05$.

The DOS given by Eq. (30) is shown in Fig. 4 for several values of the parameter α .

In the limit $\alpha = 0$, Eq. (30) reproduces the known result for class CI [6]:

$$\langle \rho_{\text{CI}}(E) \rangle = \frac{\pi}{2} \{ x [J_0^2(x) + J_1^2(x)] - J_0(x)J_1(x) \}. \quad (31)$$

A small magnetic field ($\alpha \ll 1$) changes the linear CI-class behavior $\langle \rho(E) \rangle \propto E$ to the quadratic C-class behavior $\langle \rho(E) \rangle \propto E^2$ at $x \sim \sqrt{\alpha}$:

$$\langle \rho(E) \rangle \approx \frac{\pi}{4} x \operatorname{erf}\left(\frac{x}{\sqrt{8\alpha}}\right), \quad (32)$$

see inset in Fig. 4. A similar modification of the level repulsion exponent takes place in the orthogonal-unitary crossover, Eq. (1).

In the limit of large a , one finds

$$\langle \rho(E) \rangle \approx 1 - \frac{\sin 2x}{2x} \frac{4\alpha}{\sqrt{16\alpha^2 + x^2}}, \quad (33)$$

which reduces to the C-class result (18) at $\alpha \rightarrow \infty$.

CONCLUSIONS

The purpose of this Letter was to study crossovers between superconducting symmetry classes. We have shown that the average DOS in the crossover region

between normal/superconducting symmetry classes with spin-rotation symmetry (Fig. 2) can be calculated using Efetov's supersymmetric σ -model of the orthogonal symmetry. We have obtained exact expressions (17) and (30) for the DOS in the crossover regions between the classes A–C and CI–C.

We thank D.A. Ivanov and P.M. Ostrovsky for stimulating discussions. This work was supported by the Russian Foundation for Basic Research (project no. 10-02-01180), the Dynasty Foundation, and the Russian Federal Agency of Education (contract no. NK-529P).

REFERENCES

1. M. L. Mehta, *Random Matrices* (Academic Press, Boston, 1991).
2. K. B. Efetov, *Adv. Phys.* **32**, 53 (1983).
3. K. B. Efetov, *Supersymmetry in Disorder and Chaos* (Cambridge Univ. Press, New York, 1997).
4. O. Bohigas, M. J. Giannoni, and C. Schmit, *Phys. Rev. Lett.* **52**, 1 (1984).
5. F. J. Dyson, *Commun. Math. Phys.* **19**, 235 (1970).
6. A. Altland and M. R. Zirnbauer, *Phys. Rev. B* **55**, 1142 (1997).
7. J. J. M. Verbaarschot, *Phys. Rev. Lett.* **72**, 2531 (1994).
8. A. Pandey and M. L. Mehta, *Commun. Math. Phys.* **87**, 449 (1983).
9. A. Altland, S. Iida, and K. B. Efetov, *J. Phys. A: Math. Gen.* **26**, 3545 (1993).
10. M. L. Mehta and A. Pandey, *J. Phys. A* **16**, 2655 (1983).
11. V. E. Kravtsov and M. R. Zirnbauer, *Phys. Rev. B* **46**, 4332 (1992).
12. F. Zhou, P. Charlat, B. Spivak, and B. Pannetier, *J. Low Temp. Phys.* **110**, 841 (1998).
13. M. A. Skvortsov, V. E. Kravtsov, and M. V. Feigel'man, *JETP Lett.* **68**, 84 (1998).
14. A. Altland, B. D. Simons, and D. Taras-Semchuk, *JETP Lett.* **67**, 22 (1998).
15. A. Altland, B. D. Simons, and D. Taras-Semchuk, *Adv. Phys.* **49**, 321 (2000).
16. R. Bundschuh, C. Cassanello, D. Serban, and M. R. Zirnbauer, *Phys. Rev. B* **59**, 4382 (1999).
17. P. M. Ostrovsky, M. A. Skvortsov, and M. V. Feigel'man, *Phys. Rev. Lett.* **87**, 027002 (2001).
18. C. W. J. Beenakker, *Rev. Mod. Phys.* **69**, 731 (1997).
19. W. Belzig and Yu. V. Nazarov, *Phys. Rev. B* **87**, 197006 (2001).
20. C. W. J. Beenakker, *Phys. Rev. B* **46**, 12841 (1992).
21. F. Constantinescu and H. F. de Groote, *J. Math. Phys.* **30**, 981 (1989).